

A PSEUDOVECTOR NUCLEAR HYPERFINE INTERACTION*

HARDEN M. McCONNELL†

GATES AND CRELLIN LABORATORIES OF CHEMISTRY, CALIFORNIA INSTITUTE OF TECHNOLOGY,
PASADENA 4, CALIFORNIA

Communicated by Richard M. Badger, June 18, 1958

In many studies of nuclear hyperfine interactions in polyatomic molecules and solids it has been assumed that the electron spin \mathbf{S} and nuclear spin \mathbf{I} are coupled through a symmetric dyadic \mathfrak{S} , corresponding to a hyperfine spin Hamiltonian,

$$\mathfrak{H}_{hf} = \mathbf{S} \cdot \mathfrak{S} \cdot \mathbf{I}. \quad (1)$$

When \mathfrak{S} is referred to a set of principal axes x, y, z , \mathfrak{H}_{hf} takes the familiar form,

$$\mathfrak{H}_{hf} = \mathfrak{S}_{xx} S_x I_x + \mathfrak{S}_{yy} S_y I_y + \mathfrak{S}_{zz} S_z I_z. \quad (2)$$

The purpose of the present brief note is to describe an extremely simple model that illustrates the necessity of generalizing (1) to include a pseudovector hyperfine coupling:

$$\mathfrak{H}_{hf} = \mathbf{S} \cdot \mathfrak{S} \cdot \mathbf{I} + \mathbf{V} \cdot (\mathbf{S} \times \mathbf{I}). \quad (3)$$

In equation (3), \mathbf{V} is a pseudovector fixed in the molecule or ionic complex.

The Model.—Consider a molecule containing a paramagnetic metal ion P and a magnetic nucleus X located in some other atom. For simplicity, let the molecular framework about P have an n -fold axis of symmetry that passes through P , where $n \geq 3$. This symmetry axis is parallel to the unit vector κ in Figure 1. The vector \mathbf{r} is drawn from P to X , and the direction of κ is defined so that $\kappa \cdot \mathbf{r} = r \cos \chi$ is positive. The unit vector \mathbf{h} is defined by the equation $\mathbf{r} = \kappa r \cos \chi + \mathbf{h} r \sin \chi$. In general, the Zeeman spin Hamiltonian for the coupling of an electronic spin of a paramagnetic ion to a uniform field \mathbf{H} is ¹

$$\mathfrak{H} = |\beta| \mathbf{S} \cdot \mathbf{g} \cdot \mathbf{H}, \quad (4)$$

where the spectroscopic splitting \mathbf{g} dyadic is

$$\mathbf{g} = \kappa \kappa g_{\parallel} + (\mathbf{h} \mathbf{h} + \omega \omega) g_{\perp} \quad (5)$$

and where $\omega = \kappa \times \mathbf{h}$. Assume the ion P to be far enough away from X that the field at P due to the nuclear magnetic moment $\gamma \hbar \mathbf{I}$ is uniform over the effective volume of P . The field at P due to the nuclear moment is

$$\mathbf{H}_I = \gamma \hbar \mathbf{T} \cdot \mathbf{I}, \quad (6)$$

$$\mathbf{T} = \frac{1}{r^3} [3\mathbf{r}\mathbf{r}/r^2 - \mathbf{U}] \quad (7)$$

and where \mathbf{U} is the unit dyadic. From equations (4) and (6) we obtain the hyperfine spin Hamiltonian,

$$\mathfrak{H}_{hf} = |\beta| \gamma \hbar \mathbf{S} \cdot \mathbf{g} \cdot \mathbf{T} \cdot \mathbf{I}. \quad (8)$$

The dyadic $|\beta| \gamma \hbar \mathbf{g} \cdot \mathbf{T}$ is not symmetric but can be written as the sum of symmetric and skew symmetric dyadics:

$$|\beta| \gamma \hbar \mathbf{g}, \mathbf{T} = \mathfrak{S} + \mathbf{E}. \quad (9)$$

For the problem at hand,

$$E = \frac{3}{2} \gamma \hbar |\beta| \frac{1}{r^3} (g_{\parallel} - g_{\perp}) (\cos \chi \sin \chi) (\boldsymbol{\kappa} \mathbf{h} - \mathbf{h} \boldsymbol{\kappa}). \quad (10)$$

The hyperfine coupling via the skew symmetric dyadic can also be written as the scalar product of two pseudovectors,

$$\mathfrak{S} \cdot \mathbf{E} \cdot \mathbf{I} = V \cdot (\mathbf{S} \times \mathbf{I}), \quad (11)$$

where

$$V = \frac{3}{2} \omega \gamma \hbar |\beta| \frac{1}{r^3} (g_{\parallel} - g_{\perp}) \cos \chi \sin \chi.$$

Therefore equation (3) rather than Equation (1) is the general form for the spin Hamiltonian when $g_{\parallel} \neq g_{\perp}$, and $\chi \neq 0, \pi/2$.




FIG. 1.—Pseudovector coupling between an electron spin situated at P and a nuclear spin at X .

Remarks.—In classical terms, the symmetrical dyadic interaction (1) or (2) is one which tends to orient the spins \mathbf{S} and \mathbf{I} parallel or antiparallel to one another, and along x , y , or z , depending on whether \mathfrak{S}_{xx} , \mathfrak{S}_{yy} , or \mathfrak{S}_{zz} is the largest. On the other hand, the pseudovector coupling (11) is one which tends to orient the spins \mathbf{S} and \mathbf{I} perpendicular to one another and to keep both spins in the plane of $\boldsymbol{\kappa}$ and \mathbf{h} in Figure 1. The pseudovector contribution to the hyperfine interaction can produce *first-order* effects on the hyperfine structure of paramagnetic resonance spectra. In fact, for certain parameters of our model, $3 \cos^2 \chi - 1 = 0$ and $g_{\perp} = 0$, the asymmetric nature of the coupling is dominant; i.e., the only coupling is between S_x and I_h . Comparable effects may also exist and be observable for electron-electron spin coupling.

* Sponsored by the Office of Ordnance Research, U.S. Army.

† Alfred P. Sloan Fellow.

‡ Contribution No. 2364.

¹ M. H. L. Pryce, *Proc. Phys. Soc. A*, **63**, 25, 1950.